

rpc00056

***Arabidopsis thaliana* At tom cell suspension culture**

Components

- Domestic delivery: A 50-mL plastic conical centrifuge tube, containing cell suspension
- Overseas delivery: A 250-mL plastic Erlenmeyer flask, containing cells placed on semi-solid medium

Notice

- Subculture the cells to fresh medium immediately after arrival [[Notes I](#), [II](#)].
- Do not store the cell culture in a refrigerator and a freezer.
- Maintain aseptic conditions of the cell culture, and work in a laminar flow cabinet.

Method

- Culture medium: mLS medium, 0.2 mg/L 2,4-D, pH 5.8 (medium no. 1) [[Materials III](#)]
- Culture conditions: 27°C, dark, 130 rpm [[Methods II](#)]
- Subculture: 7-day intervals [[Methods I](#)]

Citation of cell line

When results obtained by using this cell line are published in a scientific journal, it should be cited in the following manner: “*Arabidopsis thaliana* At tom cell line (rpc00056) was provided by the RIKEN BRC through the National BioResource Project of the MEXT, Japan.”

Experimental Plant Division
 RIKEN BioResource Research Center (BRC)
 Koyadai 3-1-1, Tsukuba, Ibaraki 305-0074
 Japan
 FAX: +81 29 836 9053
 E-mail: plant.brc@riken.jp
<http://epd.brc.riken.jp/en/>

Introduction

Arabidopsis At tom cell line was established from a seedling of *Arabidopsis thaliana* (L.) Heynh. accession Columbia a triple mutant that has *tom1-2* (ethyl methanesulfonate mutagenesis), *tom3-1* (ethyl methanesulfonate mutagenesis), and *thh1-1* (T-DNA insertion) mutations (Nishikiori *et al.* 2011). The At tom protoplasts show complete inhibition of tobamovirus multiplication. The At tom cells are grown in a modified Linsmaier and Skoog (mLS) medium supplemented with 0.2 mg/L 2,4-dichlorophenoxyacetic acid (2,4-D), pH 5.8. Our At tom cell culture has been maintained in the dark at 27°C with rotary shaking at 130 rpm and subcultured at 7-day intervals.

Materials

Chemicals and stock solutions

(All stock solutions are stored at 4°C)

A) MS salt mix

Murashige and Skoog Plant Salt Mixture, FUJIFILM Wako Pure Chemical Corporation (#392-00591)

B) Sucrose

C) BY2_P

KH_2PO_4 80 mg/mL

D) LS_VT_modified

Thiamine·HCl 0.4 mg/mL
myo-Inositol 40 mg/mL

E) 2,4-D (0.2 mg/mL)

2,4-D sodium monohydrate 0.236 mg/mL
(2,4-Dichlorophenoxy)acetic acid sodium salt monohydrate, Sigma-Aldrich (D6679)

F) KOH (1 N)

Glassware

A) Erlenmeyer flask (300 mL), capped with two layers of aluminum foil

B) Pipette (10 mL; large tip opening) and a bulb, sterilized by autoclaving at 121°C for 20 min

Preparation of mLS medium (medium no. 1)

1. Dissolve the following chemicals in approximately 800 mL of distilled water.

MS salt mix	1 bag (1 L)
Sucrose	30 g

2. Add following stock solutions, and fill up to approximately 950 mL with distilled water.

BY2_P	2.5 mL
LS_VT_modified	2.5 mL
2,4-D (0.2 mg/mL)	1 mL

3. Adjust the pH of the solution to 5.8 with KOH (1 N), and fill up to 1 L with distilled water.
4. Pour 70 mL of the medium into a 300-mL flask.
5. Autoclave the flask at 121°C for 20 min.

Methods

1. Leave to stand a flask containing a 7-day-old culture until the cells settle to the bottom of the flask.
2. Transfer 15 mL of the sedimented cells to 70 mL of fresh mLS medium with a pipette.
3. Incubate cell cultures on a rotary shaker at 130 rpm under the dark condition at 27°C.

Notes

- For domestic customers: We send At tom cell suspension in a 50-mL disposable conical centrifuge tube. The cells should be transferred to fresh mLS medium immediately after arrival. Transfer settled cells to Erlenmeyer flasks containing fresh liquid medium with a pipette.
- For overseas customers: We send At tom cells placed on semi-solid mLS medium in a 250-mL disposable Erlenmeyer flask. The cells should be transferred to fresh mLS medium immediately after arrival. Collect the cells from the semi-solid medium with a spatula and transfer them to Erlenmeyer flasks containing fresh liquid medium.
- In order to maintain At tom cell suspension cultures stably, it is essential to transfer an adequate amount of cells to fresh mLS medium in every subculture. The amount

of cells may vary from one lab to another, because proliferation of At tom cells is affected by culture conditions, such as a room temperature, rotation speed of a rotary shaker, and aeration condition of the culture.

- In order to obtain good aeration of a suspension culture, a silicone sponge plug may be used instead of the aluminum foil cap (*e.g.*, cap-type Silicosen; Shin-Etsu Polymer, Tokyo, Japan; <https://www.shinpoly.co.jp/en/product/product/medical/plugs.html>).
- If too many cells are subcultured to 70 mL of mLS medium, the cell culture will be overgrown after seven days. The overgrown culture shows dull yellow color. Repeated subculture of such an overgrown culture may sometimes cause sudden arrest of cell growth.

References

Nishikiori M, Mori M, Dohi K, Okamura H, Katoh E, Naito S, Meshi T, Ishikawa M (2011) A host small GTP-binding protein ARL8 plays crucial roles in tobamovirus RNA replication. PLoS Pathogen 7: e1002409. DOI: [10.1371/journal.ppat.1002409](https://doi.org/10.1371/journal.ppat.1002409)

Appendix A: Formulation of culture medium

Table A1. modified Linsmaier and Skoog medium
(medium no. 1)

Chemical	Concentration (mg/L)
KNO ₃	1900
NH ₄ NO ₃	1650
CaCl ₂ ·2H ₂ O	440
MgSO ₄ ·7H ₂ O	370
KH ₂ PO ₄	370
H ₃ BO ₃	6.2
MnSO ₄ ·4H ₂ O	22.3
ZnSO ₄ ·7H ₂ O	8.6
KI	0.83
Na ₂ MoO ₄ ·2H ₂ O	0.25
CuSO ₄ ·5H ₂ O	0.025
CoCl ₂ ·6H ₂ O	0.025
FeSO ₄ ·7H ₂ O	27.8
Na ₂ -EDTA	37.3
Thiamine·HCl	1
<i>myo</i> -Inositol	100
Sucrose	30000
2,4-D sodium monohydrate	0.236