

rpc00038

## ***Lithospermum erythrorhizon* OM callus culture**

### **Components**

- A 9-cm plastic Petri dish, containing cells placed on semi-solid medium

### **Notice**

- Subculture the cells to fresh medium immediately after arrival [[Notes I](#)].
- Do not store the cell culture in a refrigerator and a freezer.
- Maintain aseptic conditions of the cell culture, and work in a laminar flow cabinet.

### **Method**

- Culture medium: mLS medium, 1  $\mu$ M IAA, 10  $\mu$ M kinetin, 1% (w/v) agar, pH 6.3 (medium no. 33) [[Materials III](#)]
- Culture conditions: 23°C, dark [[Methods II](#)]
- Subculture: 28-day intervals [[Methods I](#)]

### **Citation of cell line**

When results obtained by using this cell line are published in a scientific journal, it should be cited in the following manner: “*Lithospermum erythrorhizon* OM cell line (rpc00038) was provided by the RIKEN BRC through the National BioResource Project of the MEXT, Japan.”

Experimental Plant Division  
RIKEN BioResource Research Center (BRC)  
Koyadai 3-1-1, Tsukuba, Ibaraki 305-0074  
Japan  
FAX: +81 29 836 9053  
E-mail: [plant.brc@riken.jp](mailto:plant.brc@riken.jp)  
<http://epd.brc.riken.jp/en/>



## Glassware and equipment

- A) Erlenmeyer flask (100 mL), capped with two layers of aluminum foil
- B) Forceps, sterilized before use

## Preparation of mLS medium (medium no. 33)

1. Dissolve the following chemicals in approximately 800 mL of distilled water.

|             |             |
|-------------|-------------|
| MS salt mix | 1 bag (1 L) |
| Sucrose     | 30 g        |

2. Add following stock solutions, and fill up to approximately 950 mL with distilled water.

|                            |        |
|----------------------------|--------|
| CuSO <sub>4</sub> (1.2 mM) | 1 mL   |
| LS_VT                      | 2.5 mL |
| IAA (1 mM)                 | 1 mL   |
| Kinetin (1 mM)             | 10 mL  |

3. Adjust the pH of the solution to 6.3 with NaOH (1 N), and fill up to 1 L with distilled water.
4. Pour 30 mL of the medium into a 100-mL flask containing 0.3 g of agar.
5. Autoclave the flask at 121°C for 20 min.

## Methods

1. Pick up an appropriate amount of callus cells from a 28-day-old culture with a forceps and place the cells onto fresh mLS medium.
2. Incubate cell cultures under the dark condition at 23°C.

## Notes

- We send OM cells on semi-solid mLS medium in a 9-cm disposable Petri dish. The cells should be subcultured to fresh mLS medium immediately after arrival.
- In order to maintain OM callus culture stably, it is essential to observe the growth of cells carefully. Because proliferation of OM cells is affected by culture conditions, such as a room temperature, aeration conditions of the culture and so on, an amount of cells transferred to fresh medium and the subculture intervals may vary from one lab to another. We usually inoculate three pieces of OM callus (about 5-mm

in diameter) on 30 mL of mLS medium in a 100-mL flask, and culture them for 28 days.

## References

Yamamoto H, Inoue K, Yazaki K (2000) Caffeic acid oligomers in *Lithospermum erythrorhizon* cell suspension cultures. *Phytochemistry* 53: 651–657. DOI: [10.1016/S0031-9422\(99\)00623-8](https://doi.org/10.1016/S0031-9422(99)00623-8)

## Appendix A: Formulation of culture medium

Table A1. modified Linsmaier and Skoog medium  
(medium no. 33)

| Chemical  | Concentration (mg/L) |
|---|----------------------|
| KNO <sub>3</sub>                                    | 1900                 |
| NH <sub>4</sub> NO <sub>3</sub>                     | 1650                 |
| CaCl <sub>2</sub> ·2H <sub>2</sub> O                | 440                  |
| MgSO <sub>4</sub> ·7H <sub>2</sub> O                | 370                  |
| KH <sub>2</sub> PO <sub>4</sub>                     | 170                  |
| H <sub>3</sub> BO <sub>3</sub>                      | 6.2                  |
| MnSO <sub>4</sub> ·4H <sub>2</sub> O                | 22.3                 |
| ZnSO <sub>4</sub> ·7H <sub>2</sub> O                | 8.6                  |
| KI  | 0.83                 |
| Na <sub>2</sub> MoO <sub>4</sub> ·2H <sub>2</sub> O | 0.25                 |
| CuSO <sub>4</sub> ·5H <sub>2</sub> O                | 0.325                |
| CoCl <sub>2</sub> ·6H <sub>2</sub> O                | 0.025                |
| FeSO <sub>4</sub> ·7H <sub>2</sub> O                | 27.8                 |
| Na <sub>2</sub> -EDTA                               | 37.3                 |
| Thiamine·HCl  | 0.4                  |
| <i>myo</i> -Inositol                                | 100                  |
| Sucrose   | 30000                |
| IAA·K   | 0.213                |
| Kinetin   | 2.15                 |
| Agar  | 10000                |